

EXPERIMENTAL AND SIMULATION STUDY OF ELECTROMAGNETIC RADIATION (EMR) EFFECT ON HUMAN THYROID TISSUE

KHITAM Y. ELWASIFE[#], M. ABUJAMI*, I. ABDELAZIZ**^{*}, M. SHABAT**

*Department of Physics, Faculty of Science, Islamic University of Gaza, Gaza Strip, Palestine,
[#]e-mail: kelwasife@iugaza.edu.ps

**Department of Biology, Faculty of Science, Islamic University of Gaza, Gaza Strip, Palestine

Abstract. This study aims to investigate the effects of non-ionizing radiation emitted from mobile phone base station on some target group of children. Their thyroid-stimulating hormone (TSH) has been investigated taking into account providing the children with possible protective olive oil supplement. The target group was composed of 120 children (6–12 years) and was divided to three sets. The first group served as control group. The second group was exposed to electromagnetic field (EMF) alone, the third group was exposed to EMF of and given 2.5 mL/day olive oil supplementation for 5 weeks. The second and the third groups lived nearby mobile phone base station (100–150 m) more than 5 years. The thyroid-stimulating hormone (TSH) was assumed. EMF exposure caused decreased in TSH. Furthermore, this work presents a simulation study of electric fields, magnetic fields, power density and specific absorption rate (SAR) distribution in human thyroid tissue. Concerning numerical modeling, the power absorption and specific absorption rate in a thyroid tissue are generally computed using FDTD methods. Results show that electromagnetic radiation (EMR) from mobile phone penetrates the thyroid tissues and attenuates rapidly to reach zero at the inner of tissue. The absorbent power and SAR show a maximum at the interface.

Key words: simulation, life tissue, cellular phone radiation, thyroid-stimulating hormone.

INTRODUCTION

According to the widespread use of mobile phones and the interactions of electromagnetic field produced from mobile phones with environment and with tissues of human beings are still under discussion and many research teams are investigating if there are any clear effects on human health, as there is still a great controversy regarding the possibility of induction of any significant physiological effects in humans by microwave radiations emitted by mobile phones. The biological effects of radiofrequency (RF) exposures from various national and

Received: May 2017;
in final form January 2018.

international expert groups had been reported [8]. The effects of electromagnetic fields induced by the Global System Mobile (GSM) communications on the thyroid-stimulating hormone in humans had been evaluated [7]. Agustíño *et al.* studied if non-ionizing radiation at 2.45 GHz may modify the expression of genes that codify heat shock proteins (HSP) in the thyroid gland [1]. They used the enzyme-linked immunosorbent assay (ELISA) technique and studied levels of HSP-90 and HSP-70. The results demonstrated that the thyroid gland is sensitive to 2.45 GHz RF and non-thermal mechanisms cause a transitory decrease in the values of HSP-90 and HSP-70, which determine cellular stress levels in the thyroid gland. Santin *et al.* investigated the possible histopathological effects of pulse modulated radiofrequency fields on the thyroid gland, using light microscopy, electron microscopy and immunohistochemical methods [6]. The aim of other researches [11–12] was to review the evidences of indirect effects on thyroid function and growth regulation, and its mechanisms. The genomic effects of electric field on normal and abnormal thyroid tissue were also reviewed, as well as no genomic, distinct molecular pathways. Simulation of human body models is used for biomedical research in a many of papers, where it is advantage to replace life human body layer by the numerical model. Biological effects of EMF are one of the areas as the numerical models with many advantages had been used [10]. On the other side, a lot of researches deal with different possibilities of numerical modeling of electromagnetic field effects on the human body which calculated the specific absorption rate distribution in human body.

In this work, we investigated the effects of radiation emitted from mobile phone base station on a target group of 30 children in control group, and 50+40=90 children in the experimental one, healthy children volunteers who had agreed to participate in this study. Their thyroid-stimulating hormone has been investigated taking into account providing the children with possible protective olive oil supplementation. We present also a simulation study of electric and magnetic fields, power density and specific absorption rate distribution in human thyroid tissue. Concerning numerical modeling, the power absorption and specific absorption rate in a thyroid tissue are effectively computed using FDTD methods.

MATERIALS AND METHODS

A total of 120 volunteer children (aging 6–12 years) participated in the experiment. They were divided into three groups: the first group (30 children) served as control group who are live outside the area of radiation, and two experimental groups, who lived, more than 5 years, in an area nearby mobile phone base station (100–150 m). So, the children where exposed to non-ionizing radiation emitted from mobile base station in Khan Younis county in Gaza Strip, Palestine,

with a power in the range of 1.4–4.7 mW/cm², which it is used in their country and measured during the experiment using power meter. The electric field is in the range of 60–130 V/m. The antenna received the signal from mobile base station at the area. The mobile system used in Gaza is GSM (Global System for Mobile) which frequency equals 900 MHz. The second group (50 children) was exposed to EMF alone, and the last group (40 children) was exposed to EMF and given 2.5 mL/day orally olive oil all over the experimental periods (5 weeks). Blood samples of 3 mL were collected. The level of TSH was measured by ELISA kits from Dia Metra, Italy [10]. Extra-virgin olive oil was taken from olive tree in Gaza which is healthy and cheap.

DATA ANALYSIS

Data were analyzed using SPSS program for windows (Statistical Package for the Social Sciences Inc. Chicago, Illinois). Mean values were compared by independent sample t-test. Significance is taken as follows: $P > 0.05$ is non-significant, $P \leq 0.05$ is significant, $P < 0.001$ is highly significant.

SIMULATION MODEL

Suppose that electromagnetic radiation with different frequencies is incident vertically upon the interface thyroid tissue [2–5]. Dielectric properties for thyroid tissue at frequency 900 MHz, shown in Table 1, were calculated by online program, Dielectric Properties of Body Tissues, from the Italian National Research Council [13].

Table 1

Dielectric properties for thyroid tissue at frequency 900 MHz

Tissue name	Conductivity σ (S/m) at 900 MHz	Relative permittivity at 900 MHz	Penetration depth (m)	Density ρ (kg/m ³)
Air	0	1	1	1.229
Thyroid	1.0385	59.684	0.040066	1100

Electromagnetic radiation research may be modeled as boundary value problems control by partial differential equations subject to initial boundary values. The spatial domain of the boundary value problem may be complicated in general. The study of a heterogeneous model for human tissue is a difficult theoretical task.

Maxwell's equations are the basic equations to simulate human life tissue by FDTD method. There have been used special models and techniques, each valid only in a limited range of frequencies or other parameters. A combination of techniques has been used to obtain specific absorption rate for different models as a function of frequency and time. The permittivity and conductivity of thyroid tissue are different according changing frequency. The electric field is assumed to propagate in the z direction with polarization at the x direction [2–5]. In addition, we assume that the fields do not vary in the x–y plane. The dielectric properties of body tissue, in general, are conceded in the complex permittivity formula as:

$$\epsilon_r^*(\omega) = \epsilon_r - j \frac{\sigma}{\omega \epsilon_0} \quad (1)$$

where ϵ_r is the relative permittivity for human tissue and changes according different frequencies, ϵ_0 is the permittivity of free space, ϵ_r^* is the complex permittivity, σ is the conductivity of the tissue and ω is the angular frequency.

According theoretical details described in the references [4, 5], in this study we choice a thyroid tissue as a model and the final equations of electric and magnetic fields in thyroid tissue are evaluated as:

$$\begin{aligned} \tilde{E}_x^{m+1/2}(\kappa) = & \frac{1 - \frac{\Delta t \cdot \sigma_{th}}{2\epsilon_0 \epsilon_{rth}}}{1 + \frac{\Delta t \cdot \sigma_{th}}{2\epsilon_0 \epsilon_{rth}}} \tilde{E}_x^{m-1/2}(\kappa) - \\ & \frac{1/2}{\epsilon_{rth} \left(1 + \frac{\Delta t \cdot \sigma_{th}}{2\epsilon_0 \epsilon_{rth}} \right)} [H_y^m(\kappa + 1/2) - H_y^m(\kappa - 1/2)] \end{aligned} \quad (2)$$

$$H_y^{m+1}(\kappa + 1/2) = H_y^m(\kappa + 1/2) - \frac{1}{\sqrt{\epsilon_0 \mu}} \frac{\Delta t}{\Delta z} [\tilde{E}_x^{m+1/2}(\kappa + 1) - \tilde{E}_x^{m+1/2}(\kappa)] \quad (3)$$

where E is the electric field, H is the magnetic field, ω is the angular frequency, ϵ_{rth} is the real relative part of the permittivity thyroid tissue, σ_{th} is the conductivity of thyroid tissue, m is the time index and κ is the spatial index, which

indexes times $t = m\Delta t$ and positions $z = \kappa\Delta z$. The time index is written as a superscript, and the spatial index is within brackets. It has been assumed that $\mu_1 = \mu_0$ where, μ_0 is vacuum permeability and μ_1 is the permeability of tissue. The free space is assumed for the exterior of the model with wave number $k_0 = \omega\sqrt{\epsilon_0\mu_0}$.

RESULTS AND DISCUSSIONS

Experimentally, the effect of electromagnetic field and the role of olive oil as protective agent on thyroid-stimulating hormone in children tissue is summarized in Table 2. Electromagnetic field exposure caused a decrease in TSH. TSH level in children tissue decreased by 28.8%, compared to the control level. The treatment of children subjected to electromagnetic field by olive oil improved the decreasing rate to 1.66%, compared to the control level. In general, according to this study, the thyroid-stimulating hormone which was subjected to electromagnetic field exposure shows a decreasing in TSH in children who are living nearby base station for five years.

Table 2

Thyroid-stimulating hormone in children tissue exposed to electromagnetic field and therapeutic action of olive oil

Parameter	Control $N = 30$	Electromagnetic field $N = 50$	Electromagnetic field + olive oil $N = 40$
TSH (mIU/L)	2.5 ± 0.2	1.78 ± 0.08	2.46 ± 0.16
%change		-28.8	-1.66
P value		< 0.01	> 0.05

In the simulation model in this study, we supposed a theoretical model in one dimensional finite difference model for predicting electric and magnetic fields in human life tissues, such as thyroid tissue, undergoing microwave heating. For cell phone radiation, the simulated radiation source is a continuous waveform of 900 MHz. The values of electrical parameters, relative permittivity and electric conductivity for this frequency, are brief in Table 1 [13]. The specific absorption rate and power density are evaluated for the human thyroid tissue using FDTD method, as the following [4]:

$$SAR = \frac{\sigma E^2}{2\rho} \quad \text{and} \quad P = \frac{\sigma E^2}{2} \quad (4)$$

where E is the electrical field intensity, σ is the electrical conductivity of the tissue, P is the power density and ρ is the mass density.

Simulations were run for global system mobile radiation of the human thyroid. FDTD method implemented to compute the distribution of electric and magnetic fields, specific absorption rate and power density in thyroid tissue. Figure 1 represent the electric field in thyroid tissue; after 150 time steps, the electric field is maximum and decrease exponentially towards the inner tissue. The magnetic field penetrate in thyroid tissue from mobile radiation is illustrated in Figure 2. It shows that the amplitude of the magnetic field at the first peak is maximum compared to the second and the third peak; it means that the effect of magnetic field in decrease fast in the inner tissue. Specific absorption rate in thyroid tissue is represented in Figure 3. We notice that the specific absorption rate value is increased in the inner tissue. In Figure 4, we represented the results of simulation of power density *vs* time steps. The maximum peak of power density is high and decreases exponentially. The power density is directly proportional with the density of the tissue, according equation (4). Figure 5 shows the conductivity *versus* frequency in thyroid tissue. The conductivity increases by increasing frequency in low frequency range, but is almost constant in high frequency range.




Fig. 1. Electric field strength in the thyroid tissue at a frequency of 900 MHz.




Fig. 2. Magnetic field strength in the thyroid tissue at a frequency of 900 MHz.




Fig. 3. Specific absorption rate in thyroid tissue at a frequency of 900 MHz.




Fig. 4. Power density in thyroid tissue at a frequency of 900 MHz.




Fig. 5. Conductivity *versus* frequency in thyroid tissue.

CONCLUSION

The response of human thyroid tissue, representing simplified model life tissue, to irradiation by plane waves produced by mobile phone, with a frequency

of 900 MHz, has been investigated. Experimentally, exposure of human thyroid tissues to electromagnetic radiation leads to a decreased in TSH; by treatment with olive oil in eating the TSH hormone is increasing after it decreased by the effect of radiation.

FDTD is used to study the distribution of the electromagnetic fields, the absorbed power, and the SAR distribution in the thyroid tissues. It is found that the fields penetrate the interface of tissue and is attenuated rapidly till they reach zero at the inner layer. Specific absorption rate has a maximum values at the interface of tissue. According to the results from the graphs, notice that the effect on the tissue of a 900 MHz radiation of the Global System Mobile is more significant at the first time steps. The dielectric properties for tissue are different according the changes in frequency.

REFERENCES

1. AGUSTIÑO, M.J.M., J.M. LEIRO, M.T.J. MORA, J.A. RODRÍGUEZ-GONZÁLEZ, F.J.J. BARREIRO, F.J. ARES-PENA, E. LÓPEZ-MARTÍN, Electromagnetic fields at 2.45 GHz trigger changes in heat shock proteins 90 and 70 without altering apoptotic activity in rat thyroid gland, *Biol. Open*, 2012, **1**(9), 831–838.
2. ELWASIFE, K.Y., Simulation of thermal global system mobile radiation in eye retina by FDTD method, *International Journal of Pure & Applied Sciences & Technology*, 2012, **10**(1), 44–50.
3. ELWASIFE, K.Y., Numerical analysis of specific absorption rate in breast fat tissue subjected to mobile phone radiation, *Journal of Emerging Trends in Computing and Information Sciences*, 2016, **7**(7), 328–331.
4. ELWASIFE, K., Power density and SAR in multi-layered life tissue at global system mobile (GSM) frequencies, *Journal of Electromagnetic Analysis and Applications*, 2011, **3**(08), 328–332.
5. ELWASIFE, K.Y., FDTD Computation of prostate tissue exposure to cellular phones radiation. Prostate, *IJET Journal*, 2012, **2**(4), 120–124.
6. ESMEKAVA, M.A., N. SEYHAN, S. ÖMEROĞLU, Pulse modulated 900 MHz radiation induces hypothyroidism and apoptosis in thyroid cells: A light, electron microscopy and immunohistochemical study, *Int. J. Radiat. Biol.*, 2010, **86**(12), 1106–1116.
7. FISHER, D.A., Physiological variations in thyroid hormones: physiological and pathophysiological considerations, *Clinical Chemistry*, 1996, **42**(1), 135–139.
8. GABRIEL, C., A. PEYMAN, E.H. GRANT, Electrical conductivity of tissue at frequencies below 1 MHz, *Physics in Medicine and Biology*, 2009, **54**(16), DOI: 10.1088/0031-9155/54/16/002.
9. GABRIEL, S., R.W. LAU, C. GABRIEL, The dielectric properties of biological tissues: II. Measurements in the frequency range 10 Hz to 20 GHz, *Physics in Medicine and Biology*, 1996, **41**(11), 2251–2269.
10. PSENAKOVA, Z., Numerical modeling of electromagnetic field effects on the human body, *Advances in Electrical and Electronic Engineering*, 2006, **319**(5), 1–2.
11. SEYED, M., H. ASADOLLAH, G. AMIR, S. RAZIEH, P. ATEFE, Alterations in TSH and thyroid hormones following mobile phone use, *Oman Med. J.*, 2009, **24**(4), 274–278.
12. SANTIN A., FURLANETTO T., Role of estrogen in thyroid function and growth regulation, *National Center for Biotechnology Information, U.S. National Library of Medicine*, doi, 2011, 875125.
13. ***Dielectric Properties of Body Tissues, <http://niremf.ifac.cnr.it/tissprop/htmlclie/htmlclie.php>.

ACCEPTED MANUSCRIPT